



fatty acid hydroxylase-associated neurodegeneration

Fatty acid hydroxylase-associated neurodegeneration (FAHN) is a progressive disorder of the nervous system (neurodegeneration) characterized by problems with movement and vision that begin during childhood or adolescence.

Changes in the way a person walks (gait) and frequent falls are usually the first noticeable signs of FAHN. Affected individuals gradually develop extreme muscle stiffness (spasticity) and exaggerated reflexes. They typically have involuntary muscle cramping (dystonia), problems with coordination and balance (ataxia), or both. The movement problems worsen over time, and some people with this condition eventually require wheelchair assistance.

People with FAHN often develop vision problems, which occur due to deterioration (atrophy) of the nerves that carry information from the eyes to the brain (the optic nerves) and difficulties with the muscles that control eye movement. Affected individuals may have a loss of sharp vision (reduced visual acuity), decreased field of vision, impaired color perception, eyes that do not look in the same direction (strabismus), rapid involuntary eye movements (nystagmus), or difficulty moving the eyes intentionally (supranuclear gaze palsy).

Speech impairment (dysarthria) also occurs in FAHN, and severely affected individuals may lose the ability to speak. People with this disorder may also have difficulty chewing or swallowing (dysphagia). In severe cases, they may develop malnutrition and require a feeding tube. The swallowing difficulties can lead to a bacterial lung infection called aspiration pneumonia, which can be life-threatening. As the disorder progresses, some affected individuals experience seizures and a decline in intellectual function.

Magnetic resonance imaging (MRI) of the brain in people with FAHN shows signs of iron accumulation, especially in an area of the brain called the globus pallidus, which is involved in regulating movement. Similar patterns of iron accumulation are seen in certain other neurological disorders such as infantile neuroaxonal dystrophy and pantothenate kinase-associated neurodegeneration. All these conditions belong to a class of disorders called neurodegeneration with brain iron accumulation (NBIA).

Frequency

FAHN is a rare disorder; only a few dozen cases have been reported.

Genetic Changes

Mutations in the *FA2H* gene cause FAHN. The *FA2H* gene provides instructions for making an enzyme called fatty acid 2-hydroxylase. This enzyme modifies fatty

acids, which are building blocks used to make fats (lipids). Specifically, fatty acid 2-hydroxylase adds a single oxygen atom to a hydrogen atom at a particular point on a fatty acid to create a 2-hydroxylated fatty acid. Certain 2-hydroxylated fatty acids are important in forming normal myelin; myelin is the protective covering that insulates nerves and ensures the rapid transmission of nerve impulses. The part of the brain and spinal cord that contains myelin is called white matter.

The *FA2H* gene mutations that cause FAHN reduce or eliminate the function of the fatty acid 2-hydroxylase enzyme. Reduction of this enzyme's function may result in abnormal myelin that is prone to deterioration (demyelination), leading to a loss of white matter (leukodystrophy). Leukodystrophy is likely involved in the development of the movement problems and other neurological abnormalities that occur in FAHN. Iron accumulation in the brain is probably also involved, although it is unclear how *FA2H* gene mutations lead to the buildup of iron.

People with *FA2H* gene mutations and some of the movement problems seen in FAHN were once classified as having a separate disorder called spastic paraplegia 35. People with mutations in this gene resulting in intellectual decline and optic nerve atrophy were said to have a disorder called *FA2H*-related leukodystrophy. However, these conditions are now generally considered to be forms of FAHN.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- dysmyelinating leukodystrophy and spastic paraparesis
- FAHN
- spastic paraplegia 35

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Spastic paraplegia 35
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C3668943/>

Other Diagnosis and Management Resources

- GeneReview: Fatty Acid Hydroxylase-Associated Neurodegeneration
<https://www.ncbi.nlm.nih.gov/books/NBK56080>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>
- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Health Topic: Degenerative Nerve Diseases
<https://medlineplus.gov/degenerativenervediseases.html>

Genetic and Rare Diseases Information Center

- Fatty acid hydroxylase-associated neurodegeneration
<https://rarediseases.info.nih.gov/diseases/10810/fatty-acid-hydroxylase-associated-neurodegeneration>

Additional NIH Resources

- National Institute of Neurological Disorders and Stroke: Neurodegeneration with Brain Iron Accumulation
<https://www.ninds.nih.gov/Disorders/All-Disorders/Neurodegeneration-Brain-Iron-Accumulation-Information-Page>

Educational Resources

- Center for Arab Genomic Studies
<http://cags.org.ae/ctga/details.aspx?id=1331>
- MalaCards: spastic paraplegia 35
http://www.malacards.org/card/spastic_paraplegia_35
- NBIAcure
<http://nbiacure.org/learn/nbia-disorders/fahn/>

- Orphanet: Neurodegeneration with brain iron accumulation
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=385
- Washington University in St. Louis Neuromuscular Disease Center
<http://neuromuscular.wustl.edu/spinal/fsp.html#spg35>

Patient Support and Advocacy Resources

- Spastic Paraplegia Foundation, Inc.
<http://sp-foundation.org/>

GeneReviews

- Fatty Acid Hydroxylase-Associated Neurodegeneration
<https://www.ncbi.nlm.nih.gov/books/NBK56080>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?term=%22fatty+acid+hydroxylase-associated+neurodegeneration%22+%5BDISEASE%5D+OR+NCT00907283+%5BID-NUMBER%5D>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28fatty+acid+hydroxylase-associated+neurodegeneration%5BTIAB%5D%29+OR+%28fa2h%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- SPASTIC PARAPLEGIA 35, AUTOSOMAL RECESSIVE
<http://omim.org/entry/612319>

Sources for This Summary

- Dick KJ, Eckhardt M, Paisán-Ruiz C, Alshehhi AA, Proukakis C, Sibtain NA, Maier H, Sharifi R, Patton MA, Bashir W, Koul R, Raeburn S, Gieselmann V, Houlden H, Crosby AH. Mutation of FA2H underlies a complicated form of hereditary spastic paraplegia (SPG35). *Hum Mutat.* 2010 Apr;31(4): E1251-60. doi: 10.1002/humu.21205.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20104589>
- Edvardson S, Hama H, Shaag A, Gomori JM, Berger I, Soffer D, Korman SH, Taustein I, Saada A, Elpeleg O. Mutations in the fatty acid 2-hydroxylase gene are associated with leukodystrophy with spastic paraparesis and dystonia. *Am J Hum Genet.* 2008 Nov;83(5):643-8. doi: 10.1016/j.ajhg.2008.10.010.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19068277>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2668027/>

- GeneReview: Fatty Acid Hydroxylase-Associated Neurodegeneration
<https://www.ncbi.nlm.nih.gov/books/NBK56080>
- Gregory A, Hayflick SJ. Genetics of neurodegeneration with brain iron accumulation. *Curr Neurol Neurosci Rep*. 2011 Jun;11(3):254-61. doi: 10.1007/s11910-011-0181-3. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21286947>
- Kruer MC, Paisán-Ruiz C, Boddaert N, Yoon MY, Hama H, Gregory A, Malandrini A, Woltjer RL, Munnich A, Gobin S, Polster BJ, Palmeri S, Edvardson S, Hardy J, Houlden H, Hayflick SJ. Defective FA2H leads to a novel form of neurodegeneration with brain iron accumulation (NBIA). *Ann Neurol*. 2010 Nov;68(5):611-8. doi: 10.1002/ana.22122.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20853438>
- Schipper HM. Neurodegeneration with brain iron accumulation - clinical syndromes and neuroimaging. *Biochim Biophys Acta*. 2012 Mar;1822(3):350-60. doi: 10.1016/j.bbadis.2011.06.016. Epub 2011 Jul 13. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21782937>
- Schneider SA, Bhatia KP. Three faces of the same gene: FA2H links neurodegeneration with brain iron accumulation, leukodystrophies, and hereditary spastic paraplegias. *Ann Neurol*. 2010 Nov;68(5):575-7. doi: 10.1002/ana.22211. Erratum in: *Ann Neurol*. 2011 Jul;70(1):187.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21031573>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/fatty-acid-hydroxylase-associated-neurodegeneration>

Reviewed: August 2015

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications

U.S. National Library of Medicine

National Institutes of Health

Department of Health & Human Services